

MNC ADCC assay on red blood cells




Figure 1

NK ADCC assay on red blood cells




Figure 2

3/16

NK ADCC: Inhibition of the ADCC by the anti-CD16: 3G8
(Tox 324 02 015)




Figure 3

4/16

Jurkat CD16 assay: activation induced by an
anti-Rhesus D and production of IL2




Figure 4

5/16

Secretion of IL2 by Jurkat CD16, induced by the
 antibodies DF5-EBV, DF5-YB20 and WinRho




Figure 5

6/16

Activation of leukocytes by anti-D antibodies




Figure 6A

Release of cytokines (IL2, IFN and TNF) by leukocytes that are antibody-activated in the presence of their target




Figure 6B

7/16

Induction of cytokine (IFN, TNF) secretion by NK cells
activated by anti-D antibodies and their target (LFB-
R297-RBC)




Figure 7A

Release of cytokines (IFN, TNF) by NK cells that are
antibody-activated in the presence of their target
(LFB-R297-RBC)




Figure 7B

8/16

Release of IL2 from Jurkat CD16, induced by anti-CD20
antibodies




Figure 8A

Secretion of IL2 by Jurkat CD16 activated by the anti-
CD20 CAT 13 or C273




Figure 8B

9/16

Release of IL2 from Jurkat CD16, induced by anti-D antibodies




Figure 9A

Secretion of IL2 by Jurkat CD16 activated by various anti-Rhesus D antibodies




Figure 9B

10/527665

Rec'd PCT/PTO 11 MAR 2005

10/16

Correlation between the ADCC assay in the presence of
Tegeline and the secretion of IL2 by Jurkat CD16.

Antibody: LFB-R297




Figure 10

11/16

IL8 secreted by MNCs in the presence or absence of target




Figure 11

12/16

Secretion of cytokines by MNCs, induced by the anti-Rhesus antibodies (deduced value without target)

Tox 324 03 062




Figure 12

13/16

Secretion of cytokines by polymorphonuclear cells,
 induced by the anti-Rhesus antibodies




Figure 13

14/16

Secretion of cytokines by NK cells, induced by the
 anti-Rhesus antibodies




Figure 14

15/16

Secretion of TNF alpha by NK cells, induced by the
anti-CD20 and anti-HLA-DR antibodies expressed in CHO
and YB2/0 (324 03 082)




Figure 15

16/16

Secretion of IFN gamma by NK cells, induced by the
 anti-CD20 and anti-HLA-DR antibodies expressed in CHO
 and YB2/0 (324 03 082)




Figure 16

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.